

CLIMATE-SMART
Agriculture
20**15**



Global Science Conference

March 16-18, 2015
Le Corum, Montpellier France

Livestock and climate change Mitigation and adaptation options for a sustainable future

Jean-François Soussana¹
and the EC FP7 'AnimalChange' consortium

www.animalchange.eu

Inra, Paris, France.

Montpellier
March 18, 2015

The AnimalChange Project

- **Funding scheme:** FP7 European Collaborative Project
- **Budget:** 9 Million euros
- **Start date:** 01/03/2011
- **Duration:** 48 months
- **Consortium:** 25 partners
- **Represented countries:** 13 European Countries, 4 African Countries, Brazil, New Zealand
- **Web site:** www.animalchange.eu



AnimalChange is supported by EU FP7 under grant agreement 266018



Dissemination component

- Stakeholder platform including key end-users
- Side events at conferences
- Development of eLearning courses
 - Mitigation and Scale, Adaptation, Equipment and Techniques for Measurement, Inventories and General Awareness for policy makers.
- Regional face-to-face training events:
 - Hungary: 27 to 31 October 2014,
 - Kenya: 1 to 5 December 2014,
 - Senegal: 12 to 16 January 2015,
 - Brazil: 9 to 13 February 2015.

The format is 3 days of training (25 Students) and 2 days of workshop



YOU ARE WELCOME TOMORROW (THURSDAY) AT THE ANIMALCHANGE SIDE-EVENT




AnimalChange is supported by EU FP7 under grant agreement 266018




AnimalChange



AnimalChange




BaU scenarios for livestock (90% increase in demand by 2050)




Livestock product price change compared to 2000 [%]

NO climate change, Middle of the road socio-economic scenario (Havlik et al.)

Increased deforestation and increased grassland cover, leading to increased GHG emissions




Reduced demand in alternative socio-economic scenarios



SSP1: more environmental
SSP3: more fragmented




Small effects of climate change on ruminants number,




Climate change impacts on land use

- Technological change and less livestock product intensive diets would lead under SSP1 to savings of 300 million hectares of forests and other natural land compared to SSP2 and to reduced emissions
- Under SSP3, because of slow productivity growth, the same amount of land would be cultivated as under SSP3, also the total livestock production is projected some 10% lower.
- Due to climate change, when CO₂ fertilization effects accounted for, the cropland is under SSP2 projected to expand by 33 million ha more than without climate change, and grassland even by 79 million ha more than without climate change.
- Given land use change, global animal numbers are not affected by climate change

Understanding and reducing uncertainties: GHG emissions (Miterrra framework)



Reducing uncertainties for soil carbon: Carbon sequestration (*NCS*) in European grasslands ($\text{g C m}^{-2} \text{ yr}^{-1}$)



$$NCS = 76 \pm 11 \text{ gC m}^{-2} \text{ yr}^{-1} \quad (P < 0.001)$$




$$NCS = (f_2 + k_N \cdot N_s) \cdot GPP + k_C \cdot L_C \quad (P < 0.001, r^2 = 0.65)$$

Critical animal stocking density for 'zero carbon' pastures

- Does C sequestration compensate for non CO₂ emissions

(CH₄, N₂O) on-site?




a. Unfertilized pasture (200 d grazing) b. Mineral N fertilized pasture (200 d grazing) c. Manure fertilized pasture (200 d grazing)



Critical herbage use efficiency = 0.20*

'Zero carbon' pastures are very extensive


Reducing uncertainties: grassland ecosystem manipulation



- Temperate: lowland sown grassland (Ireland)
- Temperate: lowland sown grassland (Switzerland)
- Temperate with future climate: upland grassland (France)
- Temperate with future climate: upland grassland (Ecotron France)
- Temperate and sub-Mediterranean: Loess grassland (Hungary)
- Semi-arid: semi-arid grassland (South Africa)
- Semi-arid: rangeland (Senegal)


Reducing uncertainties: climate change impacts on global grasslands

From 2005-2015 to 2090-2099, Orchidee (LSU/ha)




Hadgem2 Rcp 4.5

HADGEM-2





Hadgem2 Rcp 8.5

IPSL-CM5




Difference between the mean values of the aridity index (b) calculated for the years 2005-2099 and 1951-2004 with $b < 25^{\text{th}}$ percentile, as represented by two climate models and two RCPs. Red to bleu colours indicate growing aridity under future climate (and vice versa for red to brown colours).



AnimalChange



MITIGATION OPTIONS: Meta-analysis of enteric CH₄ mitigation from diet changes and additives




Meta-Gate data base



M&A option: grass-legume mixtures have increased total N yield

And increased drought tolerance in European drought experiments





- Mixtures, on average, outperform grass monocultures at majority of site
- Average N yield of mixtures is at level of legume monocultures

Tropical grazing and browsing: mitigation from secondary plant compounds

- Tropical (e.g. legumes) tannin-rich forage plants reduce methane production *in-vitro*,
- Some of the saponin-rich browsed species could also lead to reduced enteric methane emissions,
- This questions the validity of CH₄ emission factors for some of the tropical grazing/browsing systems

AnimalChange




Model & showcase farms




The model farms are virtual farms representing a livestock system in a region. These virtual farms are created combining estimated data and information based on regional production systems and national statistics.

Showcase farms, however, are real farms. Real-life data are collected on these farms. Data from showcase farms are used in calculations of model farms.

Farm AC model: modeling M&A options



Carbon fluxes



Nitrogen fluxes


M&A options: trade-offs and synergies (Dutch dairy farm example)

Mitigation Measures	Adaptation Measures			
	Water management	Fertilisation rate	Cooling of animals	Supplemental feeding
Genetic improvement in dairy cattle	No Effect	No Effect	+	No Effect
Increasing housing	+	-/+	+	-/+
Feeding maize and less grass	-	-/+	-/+	+
Legumes in the rotation	+	-	No effect	+
Replacement rate cattle	No Effect	No Effect	-/+	-/+



Note ++ is highly positive, + is positive, - is negative and – is highly negative, boxes coloured green are associated with low uncertainty, yellow is medium uncertainty and red is high uncertainty

Distribution of the adaptation and mitigation options proposed by farmers

Survey of 196 livestock farms spread over 10 countries: France, Scotland, Netherlands, Ireland, Senegal, Burkina-Faso, Kenya, South Africa, Madagascar, Brazil.




AnimalChange




Mitigation abatement cost curve for the Brazilian cerrado (2006-2030)

Marginal Cost Effectiveness
(Reals per tonne of CO₂
equivalent per year)




Index of Behavioural Barriers (IBB)



Low IBB: low uptake of M & A strategies

Behavioural barriers in Eastern Kenya



Mitigation package for Europe

Total emissions from ruminants	
Ruminants	496 Mt CO ₂ -eq.
Mixed dairy	166 Mt CO ₂ -eq.
Mitigation potential	
Fat supplementation (mixed dairy cows)	2.3 to 6.8 Mt CO ₂ -eq.
Manure management (mixed dairy)	5.3 Mt CO ₂ -eq.
Biogas production (mixed dairy)	4.4 Mt CO ₂ -eq.
Energy use efficiency (all ruminants)	23.2 Mt CO ₂ -eq.
Increase legumes in grassland (all ruminants)	14.6 Mt all systems
Grazing management (all ruminants)	0.17 Mt all systems
Total	50 to 54.5 Mt CO₂-eq Or c.a. 11%



Thank you for your attention!



AnimalChange is supported by EU FP7 under grant agreement 266018

